



Omega-3 Salmon Oil Plus: Putting “heal” in healthy!

Healing is a natural and often overlooked part of being healthy. We all assume that if we get injured we will heal. For the vast majority of us that’s exactly what happens, and without any real thought on our part. But healing and being fully healed is not necessarily the same thing. Today we know that large numbers of us live in a “nowhere zone”, never really being fully healed. Why that is so and what it means to our health is at the center of scientific investigation that holds great promise for an even healthier future for all of us...and GNLD is right there.

Decades of medical investigation has shown us that there is much more to healing than just closing a wound or mending a broken bone. Healing is also a natural and vital part of our immune function that goes on 24/7 for our entire lives. In addition to its more obvious activities, healing is involved in cleaning up the damage and debris associated with the body’s constant renewal, repair and replacement of cells. (We replace the majority of our body’s +/- 75 trillion cells every 7 years. That’s about 10 trillion cells per year or a bit more than 25 billion cells each day.) Every bump, break, bang and jolt to the body, no matter how minor, creates some damage and requires some healing. How well all of this happens has a direct bearing on our health, vitality and longevity.

Today’s science tells us that being as healthy as possible means that we have to be as “healed” as possible. The two ideas are so directly related that you can’t have one without the other. Investigating how healing happens has become one of the most exciting new frontiers of health science. Not surprisingly there is a direct connection between what you eat and how “healed” you can be.

As always GNLD is at the forefront of new understanding when it comes to nutrition and health. At a recent major scientific event, evidence from new research that used our Omega-3 Salmon Oil Plus showed the powerful healing potential of this product and the nutrients it contains. Resulting from the analysis of blood samples obtained in a human clinical trial, investigators showed that Salmon Oil Plus supported the body’s natural immune function and at the same time promoted healing!

You can read more about this study, the relationship of omega-3 fatty acids to healing and the benefits shown for Salmon Oil Plus in the attached article co-written by SAB member Dr. Arianna Carughi and Dr. Karsten Gronert, one of the world’s leading omega-3 researchers.

This is exciting and powerful new information about being as healthy as possible. It is an even more exciting and powerful demonstration of the benefits of our Omega-3 Salmon Oil Plus.

A handwritten signature in black ink, appearing to read "John R. Miller". The signature is fluid and cursive, with a long, sweeping tail on the final letter.

John R. Miller
VP Science & Technology
SAB Member

INFLAMMATION:

OMEGA-3 FATTY ACIDS AND THEIR HEALTH / DISEASE CONNECTION



ARIANNA CARUGHI

Ph.D., C.N.S.,
Nutrition Scientist,
Stanford Fellow,
GNLD SAB Member



KARSTEN GRONERT

M.S., Ph.D.
Associate Professor, Solon M. &
Pearl A. Braff Chair in
Clinical Optometric Science
Ocular Inflammation,
Vision Science Program
School of Optometry
University of California, Berkeley



MAY 18, 2010

INFLAMMATION IS ESSENTIAL TO LIFE— PART OF THE BODY'S NATURAL RESPONSE TO INJURY OR INFECTION

When in appropriate balance, inflammation comes and goes on an “as needed” basis; doing its job and then going away. However an upset balance in the level of inflammation can make it persistent or chronic and can cause progressive damage to the body and lead to a variety of diseases. Current research shows that omega-3 fatty acids are building blocks for key compounds—resolvins and protectins—that are involved in the resolution of inflammation. Experimental and clinical data provide compelling evidence that low levels of these protective omega-3 derived compounds lead to misguided and uncontrolled inflammation. It is now recognized that this may increase risk of chronic disease and mortality.

INFLAMMATION— WHAT IS IT?

Inflammation is a natural part of a healthy immune system acting as the body's basic response to injury, stress, and infection. It is essential to health and survival of all living organisms¹. It is the result of a sequence of tightly controlled, complicated, and interrelated events. It works to protect a wound against invading bacteria/viruses, remove dead tissue and cells, and repair injured or stressed tissue. By design, a healthy inflammatory response is beneficial, temporary, resolves naturally, and restores normal function of the affected tissue/organ. At any given time in life there are many points where healthy and temporary inflammation goes unnoticed in the skin, gastrointestinal tract, mouth, eyes, blood vessels, and lungs providing routine protection against potential infection, repairing minor injuries, and maintaining normal organ function.

THE PROCESS— WHAT HAPPENS?


The complex inflammatory response is driven by white blood cells (*leukocytes*) that patrol the body and possess a potent arsenal of bactericidal agents and chemical messengers that regulate inflammation, immune responses, blood vessel formation, and wound healing. Injury or infection, for a short time, will attract leukocyte to the affected area where their activity is tightly controlled to let limited and specific release of the potent chemicals that kill harmful bacteria and promote wound healing, all the while trying to limit collateral damage to the surrounding tissue.



Unregulated inflammation is now recognized as a major factor in the cause of cardiovascular diseases, asthma, arthritis, allergies, and cancer.

INFLAMMATION AND DISEASE— WHEN A GOOD THING GOES WRONG

A complex assortment of chemical messengers that include lipid signals (molecules made from specific lipids) are released to trigger inflammation and activate patrolling immune system “guardians” called leukocytes. Unfortunately, the complex regulation of healthy inflammation is in a precarious balance that can tip towards causing unwanted tissue damage and lead to abnormal or chronic inflammation^{2,3}. Poor diet or overall state of health, environmental factors, and other influences can all tip healthy inflammatory balance toward an uncontrolled and disease causing “pro-inflammatory” state. Unregulated inflammation is now recognized as a major factor in the cause of cardiovascular diseases (atherosclerosis, heart attacks, stroke), asthma, arthritis, allergies, and cancer¹. Lipid signals have long been recognized as key messengers and regulators of inflammation and thus it is not surprising that they are key targets for drugs such as corticosteroids and non-steroidal anti-inflammatory drugs (NSAID's such as aspirin and ibuprofen) that are the first treatment option for most symptoms of inflammation like pain, fever, and headaches, and inflammatory diseases like arthritis.



An imbalance of excessive omega-6 PUFA and deficient omega-3 PUFA (fish oils) consumption is associated with a higher risk of cardiovascular mortality and other inflammatory diseases.

DIETARY FATTY ACID INTAKE REGULATES INFLAMMATORY/ANTI-INFLAMMATORY BALANCE

Virtually every cell in the body can form lipid signals from key building blocks known as eicosanoids. The lipid messengers and regulators of inflammation are formed from the essential omega-6 polyunsaturated fatty acid (PUFA), arachidonic acid, while those for anti-inflammation are formed from omega-3 PUFA's. Both types of PUFA's are obtained directly through diet.

The “inflammatory” omega-6 fatty acid (arachidonic acid) or its precursor linoleic acid is the predominant PUFA in the Western Diet. The excessive omega-6 consumption in the Western diet has been linked to amplified and prolonged inflammation. Conversely, the relative lack of “anti-inflammatory” omega-3's undermines the body's ability to resolve inflammation. Thus it is not surprising that an impressive body of clinical and experimental studies have shown that an imbalance of excessive omega-6 PUFA and deficient omega-3 PUFA (fish oils) consumption is associated with a higher risk of cardiovascular mortality and other inflammatory diseases⁴. According to the American Heart Association cardiovascular disease claimed 831,272 lives in 2006, which represents 1 of every 2.9 deaths in the USA. A recent scientific meeting on “omega-3 PUFA fatty acids—recommendations for therapeutics and prevention” concluded that increasing omega-3 PUFA in tissues may lower illnesses that account for the largest diseases burden worldwide⁴.

ENDING INFLAMMATION: RESOLUTION

Traditionally, it was assumed that healthy inflammation stopped due to the fact that inflammatory signals fizzle-out and leukocytes eventually leave the injured tissue. However, research in the last 10 years has uncovered that it is a very active process controlled by many chemical signals and highly specialized pathways for the removal of spent leukocytes^{3,5,6}. A meeting of leading world scientists in the field of inflammation in 2007³ defined that healthy inflammation requires activation of so called “resolution pathways” and more importantly identified lipid signals as key messengers and regulators of this essential process. Leading laboratories around the world are now focused on developing new drugs that amplify these “resolution pathways” rather than limiting drug options to non-specific inhibition of essential inflammation and immune responses^{7,8,9}. More importantly, several recent studies have demonstrated that increased dietary intake of omega-3 PUFA amplifies these “resolution pathways”^{12,13,14}.


RESTORING BALANCE: OMEGA-3 FATTY ACIDS, RESOLVINS AND PROTECTINS

In early 2000, research teams at Harvard and Louisiana State University Health Science Center discovered that the omega-3 fatty acids eicosapentaenoic acid (EPA) and docosahexaenoic acid (DHA) are converted to potent anti-inflammatory, neuroprotective and pro-resolving lipid signals called protectins and resolvins^{7,8,10,11}. This was a major breakthrough since decades of research have failed to produce a clear mechanism for the remarkable and essential properties of dietary omega-3 fatty acids. Intense research efforts and publications in leading scientific journals in the last 5 years have established that these protective EPA and DHA signals are specifically formed during the resolution of inflammation and that their potent actions offset pro-inflammatory signals, hold-back leukocyte activation, and promote termination of inflammation in models of stroke, asthma, inflammatory bowel disease, eye diseases, diabetes, and kidney failure^{7,8,10,12,13}. More importantly, recent experimental data demonstrates that dietary omega-3 fatty acids amplify this “resolution pathway” and that their remarkable protective effects that ensure healthy inflammation are carried out through the action of EPA and DHA derived protectins and resolvins^{12,14}.

OMEGA-3 FATTY ACID SUPPLEMENTATION ATTENUATES INFLAMMATORY RESPONSE

The normal levels of these protective DHA and EPA signals in healthy humans are currently unknown. To determine if these newly discovered compounds are indeed formed in healthy individuals and if increasing tissue levels of omega-3 fatty acids restrain leukocyte activation, we collaborated with the Health Research and Studies Center in a dietary intervention study. Thirty healthy individuals were enrolled and received daily DHA and EPA supplements (460 mg DHA, 480 mg EPA) for 8 weeks. Full profiles of lipid signals in the blood and markers for the activation of leukocytes were measured. The detailed analysis of lipid signals demonstrated that activated leukocyte in all enrolled subject generates significant levels of the protective DHA lipid signals (resolvins and protectins) prior to any dietary supplementation with omega-3 fatty acids. More importantly, 8 weeks of DHA and EPA supplementation prevented leukocytes from becoming fully activated during a vascular inflammatory stimulus, even though essential inflammatory responses such as blood clotting were unaffected by the fish oil supplementation.

In summation, it has long been known that a healthy balance between omega-6 and omega-3 fatty acids was important for health protection and disease prevention. Recent experimental and clinical data have established that omega-3 fatty acid are essential for the production of lipid signals that are the key components in inflammation “resolution pathways” that ensure healthy inflammation and promote natural healing processes. This represents yet another example of the remarkable protective effects of essential omega-3 fatty acids. Experimental and clinical data provide compelling evidence that low levels of protective EPA and DHA lipid signals lead to misguided and uncontrolled inflammation and leukocyte activation. More importantly, these protective pathways directly depend on and can readily be amplified by dietary intake of DHA and EPA.



Omega-3 fatty acid are essential for the production of lipid signals that are the key components in inflammation “resolution pathways” that ensure healthy inflammation and promote natural healing processes.

REFERENCES

1. Kumar, V., Fausto, N., Abbas, A. (2004) Robbins and Cotran Pathological Basis of Disease, W.B. Saunders Company, Philadelphia
2. Martin, P., and Leibovich, S. J. (2005) Inflammatory cells during wound repair: the good, the bad and the ugly. *Trends Cell Biol*15, 599-607
3. Serhan, C. N., Brain, S. D., Buckley, C. D., Gilroy, D. W., Haslett, C., O'Neill, L. A., Perretti, M., Rossi, A. G., and Wallace, J. L. (2007) Resolution of inflammation: state of the art, definitions and terms. *FASEB J*21, 325-332
4. Hibbeln, J. R., Nieminen, L. R., Blasbalg, T. L., Riggs, J. A., and Lands, W. E. (2006) Healthy intakes of n-3 and n-6 fatty acids: estimations considering worldwide diversity. *Am J Clin Nutr*83, 1483S-1493S
5. Schwab, J. M., Chiang, N., Arita, M., and Serhan, C. N. (2007) Resolvin E1 and protectin D1 activate inflammation-resolution programmes. *Nature*447, 869-874
6. Serhan, C. N., and Savill, J. (2005) Resolution of inflammation: the beginning programs the end. *Nat Immunol*6, 1191-1197
7. Serhan, C. N., Chiang, N., and Van Dyke, T. E. (2008) Resolving inflammation: dual anti-inflammatory and pro-resolution lipid mediators. *Nat Rev Immunol*8, 349-361
8. Serhan, C. N., Yacoubian, S., and Yang, R. (2008) Anti-inflammatory and proresolving lipid mediators. *Annu Rev Pathol*3, 279-312
9. Gilroy, D. W., Lawrence, T., Perretti, M., and Rossi, A. G. (2004) Inflammatory resolution: new opportunities for drug discovery. *Nat Rev Drug Discov*3, 401-416
10. Bazan, N. G. (2006) Cell survival matters: docosahexaenoic acid signaling, neuroprotection and photoreceptors. *Trends Neurosci*29, 263-271
11. Serhan, C. N., Clish, C. B., Brannon, J., Colgan, S. P., Chiang, N., and Gronert, K. (2000) Novel functional sets of lipid-derived mediators with antiinflammatory actions generated from omega-3 fatty acids via cyclooxygenase 2-nonsteroidal antiinflammatory drugs and transcellular processing [In Process Citation]. *J Exp Med*192, 1197-1204
12. Gronert, K. (2008) Lipid autacoids in inflammation and injury responses: a matter of privilege. *Mol Interv*8, 28-35
13. Gronert, K., and Hassan, I. R. (2007) Reaping the benefits of renal protective lipid autacoids. *Drug Discovery Today: Disease Mechanisms*4, 3-10
14. Gonzalez-Periz, A., Horrillo, R., Ferre, N., Gronert, K., Dong, B., Moran-Salvador, E., Titos, E., Martinez-Clemente, M., Lopez-Parra, M., Arroyo, V., and Claria, J. (2009) Obesity-induced insulin resistance and hepatic steatosis are alleviated by omega-3 fatty acids: a role for resolvins and protectins. *FASEB J*23, 1946-1957